
## G-tube 1 hour inservice



## <u>Across</u>

3. is an infection of the skin around the tube site.

- 4. Aspirate stomach contents with a \_ to check residual.
- 7. You must have an order to \_\_\_\_\_ or
- mix medications. 8. Call the nurse \_\_\_\_\_ \_\_\_\_ if tube is
- pulled out.
- 11. Provide \_\_\_\_\_ care frequently due to NPO.
- 14. \_\_\_\_\_ tract is the continuous path from the mouth to the anus.
- 16. Call the \_\_\_\_\_ if you have difficulties.
- 20. After medication administration
- \_\_\_\_ on the MAR.
- 21. Always do \_\_\_\_\_ checks when administering medications.
- **22.** You need <u>hour of inservicing a</u> year to maintain certification.

- 24. A g-tube is placed into the \_
- 25. Get a new syringe \_
- 26. Regular \_\_\_\_\_ will help prevent clogged tubes.
- 28. \_ residual once the amount is noted.

30. Check placement by \_ \_ air and listening with a stethoscope. <u>Down</u>

- 1. Do not \_ \_\_\_\_ water or formula to warm it up.
- 2. Position the body at least at a 45 degree angle for a bolus feeding.
- 5. If you don't return the stomach
- contents you risk \_\_\_\_\_ and electrolyte imbalance.

6. When pouring liquid medications you should pour \_\_\_\_\_ from the label.

9. Someone with \_ \_\_\_\_\_ swallowing may need a g-tube.

10. When doing a treatment after a tube feed you need to \_\_\_\_\_ gloves prior. 12. J-tubes are placed into the small bowel known as the

- **13.** NPO means nothing by
- 15. Notify the nurse if you see skin
- 17. You \_\_\_\_\_ call 911 if breathing problems occur.

**18.** To prevent infection \_\_\_\_\_ your hands. 19. the syringe during a bolus

feeding to slow the flow.

23. When checking placement by injecting into the tube use at least 10cc.

27. Put on \_\_\_\_\_ before starting the tube feeding.

\_ tubing before disconnecting a 29. syringe to prevent backflow.