Name:	Date:	

Routes

	¹ s C	Α	L	Е		² N					
											3 ,
4 - 5 - 1 1 - 1			ı	6	I	E					A
⁴ T	E R A	L		°Н		В					S
H				Α		U					P
E		⁷ C		N		L		⁸ T			Π
R 9 P O	_ I Q	U	ı	D		I		R		¹⁰ D	R
Α		В		F		Z		Α		0	Α
Р		I		Е		Α		N		R	Т
E 112		С		Е		Т		S		S	I
U 5	¹² L	С		D		I		D		0	0
T 13 N T	R A P	Е	R	П	Т	0	N	Е	Α	L	N
1 0	R	N		N		N		R		Α	Р
C 2	Y	Т		G	'		•	М		Т	N
L 2	N	I			•	¹⁴ B		Α		Е	Е
E	X	М		¹⁵ L		Υ		L		R	U
V		E		0		М				Α	M
E 16S U	3 C U	Т	Α	N	Е	0	U	S		L	0
L		Е		G		U			_		N
		R				Т					ı
			1			Н					Α

Across

- **1.** What are the numbers on a syringe?
- 5. Into the digestive system.
- **9.** What can you give after medicated an animal with a solid pill?
- **13.** What is the term for the acronym IP.
- **16.** Most common route of administration, in clinic.

Down

- **2.** To turn liquid into a fine mist.
- **3.** What may force feeding an animal cause?
- 4. Term for desired effect.
- **6.** What is a better technique than leaving the food in the kennel?
- **7.** One mililiter is the same as...
- **8.** The administration that means to apply medication topically to the skin.

- **10.** Preferred sire or most subcutaneous injections.
- **11.** What gauge needle should be used for intramuscular injections?
- **12.** You can place a pill into an animal's _____ and then hold the mouth closed.
- **14.** It is easiest to give a dog medications via which route?
- **15.** Food aversion is caused by leaving food in a kennel for a _____ period of time.