


Name: \_\_\_\_\_

Date: \_\_\_\_\_

# Zoonosis



## Across

2. cysts may occur in a variety of internal organs in the human intermediate host
4. intermediate host: ctenocephalides felis, ctenocephalides canis
7. A zoonotic skin condition caused by an ectoparasitic, tunneling sarcoptiform mite
10. intermediate host: fish, frogs, and snakes
13. Transmission ingestion of eggs from soil or contaminated hands.
14. Any disease or parasite is transmissible from animal to human

15. Preparation period for Diphyllbothriasis is five to \_\_\_\_ weeks.

17. Paratenic host to toxocara canis are.

18. Almost every warm blooded animal becomes infected with \_\_\_\_.

19. transmission ingestion of raw, insufficiently cooked, or lightly pickled freshwater fish

20. first reported in farm and dairy workers

## Down

1. An ascarid that normally resides in the small intestine of the racoon.

3. Is a skin condition by the migration of hookworm larvae in the skin of the human.

5. One treatment for beef tapeworm.

6. Location is lung, occasionally other places in the body.

8. Treatment for toxocara canis.

9. Transmission ingestion of raw or partially cooked pork product or wild game.

11. ingestion of raw undercooked "Measly beef", "beef measles"

12. One treatment for Hydatid Disease.

16. the pork tapeworm, parasitizes the small intestine of human