


Name: \_\_\_\_\_

Date: \_\_\_\_\_

# Top 50 Meds



**Across**

- 2. Max Dose in 24 hours is 3-4Grams
- 5. Used in the initial treatment of MI
- 7. ACE inhibitor
- 9. Check apical heart rate for 1 full minute before administration
- 16. Proton pump inhibitor
- 18. Antiemetic
- 20. "Pain patch"
- 21. Subcutaneous anticoagulant given in the abdomen

- 23. Used for candida albicans

- 25. Rescue inhaler
- 27. Bulk forming laxative
- 28. Thyroid supplement

**Down**

- 1. Beta-blocker
- 3. Antihistamine used for seasonal allergies
- 4. Therapeutic INR 2-3
- 6. Used for BPH
- 8. Muscle relaxant
- 10. Decongestant
- 11. NSAID

- 12. Often used for anxiety, but can also be used to stop seizure activity
- 13. Anti-Parkinson med
- 14. Med used for glucose control in type II diabetics
- 15. Combination of 2 types of pain meds-hydrocodone and acetaminophen
- 17. Mainly used for nerve pain
- 19. Sedating antihistamine
- 22. Sulfonamide antibiotic
- 24. Used to lower cholesterol
- 26. Long acting insulin