
Name:	Date:
-------	-------

B Positive: A Blood Transfusion Crossword Puzzle



<u>Across</u>

- 2. Type of transfusion reaction in which symptoms develop immediately and include facial flushing, chills, headache, and lower back pain
- **5.** Swelling, redness, and pain present at the patient's IV site during a blood transfusion indicates that this has occurred
- **10.** Patient assessment data that should be obtained by the nurse prior to beginning a blood transfusion
- 11. Two of these people are responsible for comparing and validating blood product information prior to a transfusion

- **12.** The presence of clumping or gas bubbles in a blood product may indicated that this has occurred; if found, blood should be returned to blood bank
- **14.** Smallest size peripheral venous catheter that should be used for blood product administration
- **15.** Signs of flushing, dyspnea, itching, hives, or a rash may indicate that this is occurring

Down

- 1. Time period during a blood transfusion where the most intense monitoring is required
- **3.** First action a nurse should take with regard to the blood transfusion if a transfusion reaction is suspected

- **4.** Maximum time to administer a blood product to reduce the chance of bacterial contamination
- **6.** The only solution that can be administered with blood products
- **7.** An identifier in addition to a patient's full name that can be used to confirm a patient's identify prior to transfusing blood
- **8.** Device found in a blood administration IV set that removes the particulate material formed during the storage of blood
- **9.** Type of IV administration set used to administer blood
- **13.** True of False: A medical order is required for transfusion of a blood product.