


Name: _____

Date: _____

Newborn Drug Abuse



Across

7. These anxiety-relieving drugs include Valium®, Xanax®, Klonopin® and Ativan®, and should be avoided during pregnancy if possible.

10. FAS

11. NAS

13. Drug used to treat opioid addiction

14. A score is given for each sign at one minute and five minutes after the birth. If there are problems with the baby an additional score is given at 10 minutes. A score of 7-10 is considered normal, while 4-7 might require some resuscitation measures, and a baby with an APGAR Score of 3 and below requires immediate resuscitation.

15. Substances and medications that are non-prescription and not obtained over-the-counter from a pharmacy

16. Dark greenish waste products that accumulate in the bowel during fetal life and are eliminated shortly after birth. Often tested to determine any drug use by the mother while pregnant.

Down

1. is a prescription medication that can help treat heroin addiction and dependence on other opiates.

2. The time period in weeks from conception to delivery.

3. NICU

4. Type of drug used to relieve pain. These addictive drugs are also called narcotics. Examples include: morphine, codeine, fentanyl, oxycodone, heroin, and hydromorphone. Opiates are naturally occurring narcotics whereas opioids include synthetic narcotics.

5. A yellow coloration of the skin and eyes caused by increased amounts of bilirubin in the blood. Bilirubin is a break-down product of red blood cells; it is processed and excreted by the liver. Treatments for jaundice include phototherapy ("bili-lights") and (rarely) exchange transfusion

6. Unborn baby

8. A tool to evaluate NAS, that measures the degree to which a neonate is experiencing symptoms of withdrawal due to maternal substance abuse.

9. The 6 week period following the birth of a baby

12. Birth to 28 day old infant