
Name:	Date:

Exploring Data



Across

- 1. The difference between the first and third quartiles
- 4. Shape where the longer tail stretches to the left
- 7. Value more than 1.5 times the IQR below Q1 or above Q3
- $\pmb{8}.$ Variable that describes data using words or numerals as labels
- 9. Tells how many standard deviations a value is from the mean $% \left(1\right) =\left(1\right) +\left(1\right) +\left($
- **15.** Variable that describes data using numbers as numerical values
- 20. Difference between the maximum and minimum value
 21. Found by summing all the data values and dividing by the count
- 23. Minimum, 1st quartile, Median, 3rd quartile, Maximum
- 25. Shape where the longer tail stretches to the right
- **27.** Uses adjacent bars to show the distribution of values in a quantitative variable, where each bar represents the number of values falling in an interval
- **30.** Reveals single vs. multiple modes and symmetric vs. skewness
- **31.** Type of display that shows quantitative data values in a way that shows the shape of the distribution in addition to individual data values

- $\textbf{36.} \ \ \textbf{Distribution of a variable when considering only a smaller group of individuals}$
- 37. In a Normal Model, about 68% of the values fall within 1 standard deviation of the mean, about 95% within 2 standard deviations, and about 99.7% within 3 standard deviations
- 39. In a two-way table, the distribution of either variable
- alone
 40. The value with a quarter of the data above it
- Down
- 2. The number that falls above a given % of the data
 3. Displays the 5-number summary as a central box with the whiskers that extend to the non-outlying data values
- 5. Type of Normal model with mean 0 and standard deviation 1
- 6. Distribution with two modes
- 10. The value with a quarter of the data below it
- 11. The square root of the variance
- **12.** Table that lists the categories of a variable and gives the proportion of observations for each category
- 13. The value found by subtracting the mean and dividing by the standard deviation
- 14. Shows bars divided proportionally into segments corresponding to the percentage in each group
- **16.** Shows how a "whole" divides into categories by showing a wedge of a circle whose area corresponds to the proportion in each category

- ${\bf 18.}\ {\bf The}\ possible\ values\ of\ the\ variable\ and\ the\ relative\ frequency\ of\ each\ value$
- 19. The middle value of a distribution with half the data above and half below it
- 22. Distributions with more than two modes
- 24. A distribution roughly flat in shape
- **26.** Numerical attribute of a population
- 28. Shape where the two halves on either side of the center look approximately like mirror images of each other
- 29. Having one mode
- 32. Graphs a dot for each case against a single axis
- 33. Shows a bar representing the count of each category in a categorical variable
- 34. Numerical attribute of a set of data
- **35.** A numerical summary of how tightly the values are clustered around the "center"
- **38.** When a distribution is not symmetric and one tail stretches out farther than the other