
Name:	Date:
-------	-------

Hip and Pelvis



Across

- 1. A painful injury caused by a direct blow to the hip, resulting in ecchymosis, tenderness, and swelling; also known as a hip pointer
- 7. The portion of the pelvis attached to the pubis in front and the ilium laterally in the back; bears the weight of the body when sitting.
- **8.** The portion of the vertebral column between the lumbar vertebrae and the coccyx; composed of five fused vertebrae.

- **9.** A space in the pelvis through which the sciatic nerve travels to the legs.
- **11.** The bone in the pelvis to the front and below the bladder.
- **14.** The upper ridge of the ilium.
- 2. A muscle group that aids in adduction of the hip; consists of the adductor longus, adductor brevis, and adductor magnus muscles.
- **3.** The broad, slightly concave inner surface of the ilium.
- **4.** The center of the pubis where the two sides of the pubis are fused together.

- **5.** The large openings in the ischium through which blood vessels and nerves pass to the legs.
- **6.** A muscle group that aids in hip movement; consists of the biceps femoris, semitendinosus, and semimembranosus muscles.
- **10.** A muscle group that aids in flexion of the hip; consists of the iliopsoas, sartorius, pectineus, and rectus femoris muscles.
- 12. The tailbone.
- **13.** A broad, flared bone that makes up the upper and lateral sections of the pelvis.