
Name:	Date:
-------	-------

## Cardiac Emergencies/Unresponsive Choking



## Across

- **2.** Use weight from here when giving compression ( ) ( )
- **4.** With an obstructed airway this is what you do after compressions (and before rescue breaths)
- **6.** Pain in the chest that comes and goes at different times
- **7.** Number of compressions for an adult, child and infant
- 9. A condition in which the heart has stopped beating is called
- **10.** Important care for a victim of a heart attack

- **14.** Most people die of a heart attack with ( ) hours
- **15.** Compress the chest at the rate of ( ) ( ) compressions per minute
- **16.** A reason to stop CPR is because another ( ) person arrives
- **18.** First step when using the AED ( ) ( ) ( )
- **19.** Give two of these after thirty compressions ( ) ( )

## Down

**1.** Stands for cardiopulmonary resuscitation

- **3.** When putting pressure on the chest during CPR these are called ( )
- **5.** Use two of these when giving compressions on an infant
- **8.** The most common symptom of a heart attack ( ) ( )
- 11. Placement (body part) of the heel of the hand when giving CPR
- **12.** If breath does not go in, ( ) and try again
- **13.** Place heel of hand in the ( ) of the sternum when giving compressions on an adult
- **17.** A lifesaving device that delivers an electric shock