
Name:	Date:	Period:
-------	-------	---------

## Drugs Used In Mental Health Disorders



## Across

- **5.** All except \_\_\_\_ are used to treat anxiety from alcohol withdrawl
- 7. Typical and Atypical agents are associated with which drug therapy?
- **9.** Drug of choice for alcohol withdrawl
- **10.** Actions: Inhibit reuptake and destruction of serotonin and norepinephrine from synaptic cleft
- **12.** Used to treat acute mania associated with bipolar disorder
- **13.** Common adverse effects: Sedation, Lethargy Serious Adverse effects: Slurred speech, Dizziness **14.** CNS depressants, reduce anxiety
- by stimulating the neurotramitter

- **18.** Block metabolic destruction of norepinephrine, dopamine, and serotonin neurotransmitters
- **19.** There are 5 types of antidepressants:Miscellaneous agents, MAOI's, SSRI's SNRI's & \_\_\_\_\_?
- **20.** Most common cause of nonadherence
- **21.** Phobic disorders often lead to exaggerated \_\_\_\_\_\_ behaviors and leave their victims functionally disabled.

## Down

- 1. Drug of choice for treating OCD?
  2. Prolong action of norepinephrine, dopamine, and serotonin by blocking reuptake
- 3. When taking MAOIs patient shouldn't ingest meals with a high

- **4.** Serious adverse effect of MAOIs? **6.** Used often for mild anxiety: NOT
- addictive

  effects can be
- **8.** \_\_\_\_\_ effects can be minimized by administering drugs at bedtime.
- **11.** Celexa, Lexapro, Prozac, Luvox, Paxil, Zoloft are ALL \_\_\_\_\_?
- **15.** Nueroleptic malignant syndrome is potentially
- **16.** Sedative-hypnotic; High potential for abuse
- 17. Partial Serotonin and Dopamine Agonist, Short-term relief symptoms of anxiety