
Name:	Date:	Period:	

Drugs Used In Mental Health Disorders



Across

- 5. All except ____ are used to treat anxiety from alcohol withdrawl7. Typical and Atypical agents are
- associated with which drug therapy?
- **9.** Drug of choice for alcohol withdrawl
- **10.** Actions: Inhibit reuptake and destruction of serotonin and norepinephrine from synaptic cleft
- 12. Used to treat acute mania associated with bipolar disorder
- **13.** Common adverse effects: Sedation, Lethargy Serious Adverse effects: Slurred speech, Dizziness
- **14.** CNS depressants, reduce anxiety by stimulating the neurotramitter

- **18.** Block metabolic destruction of norepinephrine, dopamine, and serotonin neurotransmitters
- **19.** There are 5 types of antidepressants:Miscellaneous agents, MAOI's, SSRI's SNRI's & _____?
- **20.** Most common cause of nonadherence
- **21.** Phobic disorders often lead to exaggerated ______ behaviors and leave their victims functionally disabled.

<u>Down</u>

- Drug of choice for treating OCD?
 Prolong action of norepinephrine, dopamine, and serotonin by blocking reuptake
- **3.** When taking MAOIs patient shouldn't ingest meals with a high

- 4. Serious adverse effect of MAOIs?
- **6.** Used often for mild anxiety: NOT addictive **8.** effects can be
- **8.** _____ effects can be minimized by administering drugs at bedtime.
- 11. Celexa, Lexapro, Prozac, Luvox, Paxil, Zoloft are ALL _____?
- **15.** Nueroleptic malignant syndrome is potentially
- **16.** Sedative-hypnotic; High potential for abuse
- 17. Partial Serotonin and Dopamine Agonist, Short-term relief symptoms of anxiety