


Name: \_\_\_\_\_

Date: \_\_\_\_\_

# Shaken Baby Syndrome



## Across

1. Fewer than \_\_\_\_\_ % of infants who survive have normal development.
5. Thirty-five percent of victims of SBS are completely \_\_\_\_\_ on their caregivers.
8. 50%-75% of \_\_\_\_\_ and adults surveyed did not know shaking an infant is dangerous.
10. Violently shaking an infant can cause injury after as few as \_\_\_\_\_ seconds.
11. Symptoms of SBS can start \_\_\_\_\_ or peek within 4-6 hours.
12. Injury occurs when shaking an infant results in force that \_\_\_\_\_ the bridging veins between the dura mater and the brain substance.

14. The mortality rate with SBS is approximately \_\_\_\_\_ %.

15. It is \_\_\_\_\_ ok to shake an infant.

16. \_\_\_\_\_ is the soft spot on an infant's head.

18. There are 1200 to 1600 reported cases of SBS in the \_\_\_\_\_ each year.

19. A long-term affect of SBS is \_\_\_\_\_.

20. The degree of damage depends on the amount and \_\_\_\_\_ of shaking that child endured.

## Down

2. Stepfathers and \_\_\_\_\_ boyfriends cause 60% of SBS cases each year.

3. SBS is a form of \_\_\_\_\_.

4. \_\_\_\_\_ is caused from violently shaking an infant.

6. SBS causes more \_\_\_\_\_ than a fall from a 3 story building.

7. Mothers are \_\_\_\_\_ in 12.6% of cases.

9. \_\_\_\_\_ is a long-term affect of SBS.

13. SBS is most common in children under \_\_\_\_\_ years old.

17. Children up to age \_\_\_\_\_ can be injured by violent shaking.