


# Shaken Baby Syndrome



**Across**

- 1. Fewer than \_\_\_\_\_ % of infants who survive have normal development.
- 5. Thirty-five percent of victims of SBS are completely \_\_\_\_\_ on their caregivers.
- 8. 50%-75% of \_\_\_\_\_ and adults surveyed did not know shaking an infant is dangerous
- 10. Violently shaking an infant can cause injury after as few as \_\_\_\_\_ seconds.
- 11. Symptoms of SBS can start \_\_\_\_\_ or peek within 4-6 hours.
- 12. Injury occurs when shaking an infant results in force that \_\_\_\_\_ the bridging veins between the dura mater and the brain substance.

- 14. The mortality rate with SBS is approximately \_\_\_\_\_%
- 15. It is \_\_\_\_\_ ok to shake an infant.
- 16. \_\_\_\_\_ is the soft spot on an infants head.
- 18. There are 1200 to 1600 reported cases of SBS in the \_\_\_\_\_ each year.
- 19. A long-term affect of SBS is \_\_\_\_\_.
- 20. The degree of damage depends on the amount and \_\_\_\_\_ of shaking that child endured.

**Down**

- 2. Stepfathers and \_\_\_\_\_ boyfriends cause 60% of SBS cases each year.

- 3. SBS ia a form of \_\_\_\_\_.
- 4. \_\_\_\_\_ is caused from violently shaking an infant.
- 6. SBS causes more \_\_\_\_\_ then a fall from a 3 story building.
- 7. Mothers are \_\_\_\_\_ in 12.6% of cases.
- 9. \_\_\_\_\_ is a long-term affect of SBS.
- 13. SBS is most common in children under \_\_\_\_\_ years old.
- 17. Children up to age \_\_\_\_\_ can be injured by violent shaking.