
IBS/IBD/GERD/PUD



Across

1. excessive fat in the stools due to malabsorption

3. generic of Uceris, used to help treat ulcerative colitis

4. Blood in stool

5. ADR of anti-histamine's, dizziness, spinning room

6. Generic name of Tagamet that has many drug interactions

8. IBS-C in women, OI-Constipation

11. Brand only, PPI

18. Side effect of GERD, ADR of Imodium

22. _____ tenderness, indicates severe to fulminant UC

23. Go to antacid, active ingredient in TUMS

25. When there are drug interactions with antacid use, separate antacid by ____ hours

28. Gives greatest symptom relief for GERD + highest healing rates

31. Malignant gastrinomas occur in 30-50% of patients with this syndrome

32. Not recommended for acid reflux, dosed QID, forms viscous adhesive on ulcer crater

33. Omeprazole and sodium bicarb

34. 250/500 mg capsules that are effective in entire GI tract

35. Weakening of bone, extra intestinal manifestation

36. used for IBS-D and hits Mu receptors

37. Food that decreases LES pressure

Down

2. A dangerous complication of GI ulcers

7. antidiarrheal that is over the counter

9. use of loperamide may cause this condition

10. classic symptom of PUD

12. Protective factor for PUD

13. Medication that is an irritant to mucosa in GERD

14. Initial treatment for ulcerative colitis

15. Prokinetic agent, increases LES pressure

16. Generic name for Sandostatin (inhibits gastric acid and releases gastrin)

17. An independent risk factor for stress-related mucosal bleeding

19. Barrier to effective treatment of H. pylori infection

20. emergency use in IBS-C and CIC

21. Mucus secreting cells in intestinal lining

24. lifestyle modification for nocturnal symptoms of GERD

26. PPI's are dosed 30-60 mins before which meal?

27. Drug used for prevention of NSAID ulcers, but contraindicated in pregnancy

29. ___ weeks of PPI is the choice for symptomatic relief and healing of erosive esophagitis

30. Food may relieve the pain when dealing with a ____ ulcer