


# New Hire Infection Control



## Across

**2.** Alcohol based hand sanitizers are acceptable in many clinical situations except for after providing care to a patient with C.

**4.** Gloves, gowns, face shields or masks, eye protection, surgical caps, and shoe covers are all examples of

**6.** Type of infection spread by contact with patient or items and surfaces in the patients environment

**7.** Isolation gowns should be removed \_\_\_\_\_ leaving the patient care area

**8.** Gloves should not be worn in a \_\_\_\_\_ unless you are carrying something soiled or dirty to another location

**10.** All \_\_\_\_\_, no matter where they come from, should be considered contaminated

**12.** When should you wash your hands after removing gloves?

**13.** The most important intervention to prevent hospital associated infections is

## Down

**1.** Type of isolation spread by oral/nasal secretions

**3.** When removing a gown, rip/pull if from your body being sure not to \_\_\_\_\_ your clothing

**5.** Who is responsible for preventing Hospital Associated infections?

**9.** Always wash hands \_\_\_\_\_ glove removal

**11.** Type of infection spread by airborne particles from oral/nasal secretions that float on air currents

## Word Bank

after

Droplet

Airborne

Immediately PPE

before

body fluids

Contact

Handwashing Diff

Everyone

Contaminate

hallway